



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

- ∴ by (1) A's portion = $\frac{1}{3}(400+000+200) = 200$,
 " (2) B's portion = $\frac{1}{3}(000+300+000) = 100$,
 " (3) C's portion = $\frac{1}{3}(000+100+200) = \underline{100}$.
 Net estate (*d*) = \$400.
-

PROBLEMS.

250. By L. REGAN, *Boonsboro, Iowa*.—Divide the line *AB*, geometrically, into three parts that shall be in harmonic proportion.

251. By DR. H. EGGERS, *Milwaukee, Wisconsin*.—Show that when the two lines, which bisect two angles of a triangle, are equal the triangle is isosceles.

252. By W. E. HEAL, *Wheeling, Ind.*.—If the roots of a given cubic equation be not *real* and *positive* show that the equation can be transformed into another, of the same degree, in which *all* the roots are real and positive.

253. By PROF. E. J. EDMUNDS, *New Orleans, La.*.—If $f(x)$ be a function whose roots are all real, show that the differential of the second order of that function has all its roots imaginary.

254. By ARTEMAS MARTIN, M. A., *Erie, Pa.*.—Integrate

$$dI = xE(e, x)dx,$$

where $E(e, x)$ denotes an elliptic arc, eccentricity e and abscissa x .

255. By PROF. JOHNSON.—1. In a triangle ABC the angle $A = \varphi + \alpha$, $B = 2\varphi$; supposing AB to remain fixed, while φ varies, it is required to find the rectangular equation to the locus of C , and the equations to the asymptotes.

2. With the same data as above, taking A as the origin, and AB as the axis of x ; let $\alpha = 45^\circ$, and find the envelop of a straight line which passes through C and makes an angle $4\varphi + 90^\circ$ with Ax .

256. By PROF. H. T. EDDY, *Cincinnati, Ohio*.—If the given quantities x_1, x_2, x_3, x_4 have the probable errors r_1, r_2, r_3, r_4 , respectively, find the probable error r of the quantity x when $x_1 : x_2 :: x_3 + x : x_4 + x$.

257. By E. B. SEITZ, *Greenville, Ohio*.—Within a triangle ABC , determine a point P , such that $m.PA + n.PB + r.PC$ shall be a minimum, m, n, r being constants.

258. By J. M. ARNOLD, *Boston, Mass.*—The Balance of a watch is slightly heavier on one side than on the other. When the watch is placed vertically with the pendant up, the heavy point is at the lowest side when the balance is at rest.

If the watch be started with a small arc of vibration the rate will be much faster, in the vertical position, with the pendant up than with the pendant down. If the motive power be increased so that the arc of vibration becomes large, say one and a half revolutions or 270° each side the point of rest, it will be found to go faster, in the vertical position, pendant down than it does with the pendant up.

What arc of vibration must be given the balance, so that its rate shall be the same in any vertical position?

QUERY 1. By PROF. HALL.—When we descend below the surface of the Earth, does the Earth's attractive force increase or diminish?

QUERY 2. By MR. KUMMELL.—Let $\cos^2\varphi$ be the probability of a single event, hence $\sin^2\varphi$ that of its non-occurrence. Supposing m simultaneous events of either kind, then we have for the mean square of the difference in number, after an infinite number of trials, the expression:

$$\epsilon_m^2 = \cos^{2m}\varphi(m)^2 + \frac{m}{1} \cos^{2m-2}\varphi \sin^2\varphi(m-2)^2 + \frac{m(m-1)}{1 \cdot 2} \cos^{2m-4}\varphi \sin^4\varphi(m-4)^2 + \dots \\ + \frac{m(m-1)}{1 \cdot 2} \dots \frac{m-n+1}{n} \cos^{2m-2n}\varphi \sin^{2n}\varphi(m-2n)^2 + \dots \sin^{2m}\varphi(-m)^2 = m(\sin^2 2\varphi + m \cos^2 2\varphi)$$

Since $m(\cos^2\varphi - \sin^2\varphi) = m \cos 2\varphi$ is the most probable difference in the number of the two kinds of events, we have $\sin 2\varphi/\sqrt{m}$ for the mean error with respect to the most probable difference. How is this proved otherwise than by induction?

PUBLICATIONS RECEIVED.

Physical and Mathematical Principles of the Nebular Theory; 10 pages, 8vo:

The Origin of the Power which Causes the Stellar Radiations; 10 pages, 8vo:

Elements of Sideral Astronomy; 7 pages, 8vo: By JACOB ENNIS, A. M., Professor of Natural Science in the State Normal School, Shippensburg, Pa.

ERRATA.

On page 47, line 5, insert $\frac{1}{2}$ before the sign of integration.

“ “ “ 7, multiply the term under the sign of integration by $d\phi$.

“ “ “ 9, for $n-3$, read $n-2$.